

Your  
Hospital's  
Logo  
Here

# ADDENDUM APS ORDERS

Medications Changes,  
Miscellaneous Orders &  
Orders to Discontinue Acute  
Pain Management

DATE: \_\_\_\_\_ TIME: \_\_\_\_\_ ( Military Time )

## PATIENT IDENTIFICATION

### MEDICATION CHANGES:

Discontinue the previous continuous epidural solution orders. Administer the following via the epidural catheter

**Date Ordered**

- \_\_\_\_\_ (1) Levo-bupivacaine (Chirocaine) 0.125% (1/8th) + Dilaudid 20 mcg/ml at \_\_\_\_\_ ml/hr
- \_\_\_\_\_ (2) Levo-bupivacaine (Chirocaine) 0.1% (1/10th) + Dilaudid 20 mcg/ml at \_\_\_\_\_ ml/hr
- \_\_\_\_\_ (3) Levo-bupivacaine (Chirocaine) 0.0625% (1/16th) + Dilaudid 20 mcg/ml at \_\_\_\_\_ ml/hr
- \_\_\_\_\_ (4) Levo-bupivacaine (Chirocaine) 0.125% (1/8th) at \_\_\_\_\_ ml/hr
- \_\_\_\_\_ (5) Ropivacaine (Naropin) 0.2% (1/5th) + Dilaudid 20 mcg/ml at \_\_\_\_\_ ml/hr
- \_\_\_\_\_ (6) Ropivacaine (Naropin) 0.1% (1/10th)+ Dilaudid 20 mcg/ml at \_\_\_\_\_ ml/hr
- \_\_\_\_\_ (7) Ropivacaine (Naropin) 0.2% (1/5th) \_\_\_\_\_ ml/hr
- \_\_\_\_\_ (8) Other \_\_\_\_\_

### TREATMENT OF BREAKTHROUGH PAIN:

For pain score of > 4 while at rest, increase infusion rate at \_\_\_\_\_ ml per \_\_\_\_\_ as needed, not to exceed \_\_\_\_\_ ml/hr.

### MISCELLANEOUS ORDERS:

1. \_\_\_\_\_
2. \_\_\_\_\_
3. \_\_\_\_\_
4. \_\_\_\_\_

### DISCONTINUATION OF EPIDURAL INFUSION:

Discontinue continuous epidural infusion at \_\_\_\_\_  
( Military Time )

Continue APS monitoring until \_\_\_\_\_ as per protocol below:  
( Military Time )

- 24 hours after last bolus dose of Duramorph
- 4 hours after discontinuing a continuous Duramorph infusion
- 4 hours after last Fentanyl bolus dose and/or local anesthetic
- 2 hours after discontinuing a Fentanyl infusion and/or local anesthetic

**TIME NOTED** Nurse's Signature \_\_\_\_\_ Doctor \_\_\_\_\_, M.D.